

Sedimentation Pond and Plant Filter Strip in Mitigating RIFA Insecticide Runoff at el modeno Gardens



Dave Kim, Johanna Walters, Kean S. Goh
Department of Pesticide Regulation
John Kabashima, Darren Haver
UC Cooperative Extension
Toby Mancini
el modeno Gardens
August 15, 2001



Outline

- Introduction
- Runoff Monitoring
- Runoff Mitigation
- Conclusions







The Insecticides






- Bifenthrin (Talstar®)
- Fenoxy carb (Award®)
- Hydramethylnon (Amdro®)
- Pyriproxyfen (Distance®)
- Chlorpyrifos (Dursban®)
- Diazinon
- Malathion
- Dimethoate

Insecticide Factors

- Vapor Pressure (mm Hg)
- Water Solubility (ppm, mg/L)
- Soil Adsorption Coefficient (Koc)
- Field Dissipation ($t_{1/2}$, days)
















Prevention & Mitigation

Insecticide & Application
Irrigation Management
Water Recovery
Holding Pond
Buffer Strip







Conclusions

Runoff

- Bifenthrin, diazinon, malathion
- Bifenthrin high >3 ppb, currently <1 ppb
- Diazinon high >8 ppb, currently trace
- Peaks at rain runoff
- Toxic to *C. dubia*

Mitigation

Reduction in Water Concentration

- **Bifenthrin 52%**
- **Chlorpyrifos 61%**
- **Diazinon -13%**
- **Malathion 31%**

Sediments

- Bifenthrin
 - Increased through time
 - 200 – 1400 ppb
 - Increased downstream basin
 - 150 – 250 ppb
 - 800 – 1200 ppb
- Chlorpyrifos <80 ppb
- Diazinon none detected



Acknowledgments

California Dept of Food and Agriculture
Center for Analytical Chemistry
UC Cooperative Extension
el modeno Gardens
Department of Fish and Game
Orange County Dept of Agriculture
Regional Water Quality Control Board

www.cdpr.ca.gov/docs/rifa